



Performance Outcomes: Measure & Metric Details

Adherence to Antipsychotic Medications for People with Schizophrenia

Numerator: Number of people who remained on an antipsychotic medication for at least 80% of their treatment period

Denominator: Number of people ages 19 to 64 with Schizophrenia who were dispensed at least 2 antipsychotic medications during the MY

Adult Access to Preventative or Ambulatory Care - 20 to 44

Numerator: Number of adults who had an ambulatory or preventative care visit during MY

Denominator: Number of adults ages 20-44 as of June 30 MY

Adult Access to Preventative or Ambulatory Care - 45 to 64

Numerator: Number of adults who had an ambulatory or preventative care visit during MY

Denominator: Number of adults ages 45-64 as of June 30 MY

Adult Access to Preventative or Ambulatory Care - 65 and older

Numerator: Number of adults who had an ambulatory or preventative care visit during MY

Denominator: Number of adults ages 65 and older as of June 30 MY

Antidepressant Medication Management - Effective Acute Phase Treatment

Numerator: Number of people who remained on antidepressant medication during the entire 12-week acute treatment phase

Denominator: Number of people 18 and older who were diagnosed with depression and treated with antidepressant medication

Antidepressant Medication Management - Effective Continuation Phase Treatment

Numerator: Number of people who remained on antidepressant medication for at least 6 months

Denominator: Number of people 18 and older who were diagnosed with depression and treated with antidepressant medication

Aspirin Use

Numerator: Number of respondents who are currently taking aspirin daily or every other day

Denominator: Number of respondents who are men, ages 46 to 65 years with at least one cardiovascular risk factor; men, ages 66 to 79, regardless of risk factors and women ages 56 to 79 with at least 2 cardiovascular risk factors

Cardiovascular Monitoring for people with Cardiovascular Disease and Schizophrenia

Numerator: Number of people who had a LDL-C test during the Measurement Year

Denominator: Number of people ages 18 to 64 years with schizophrenia and cardiovascular disease

Care Coordination (Q13, 17 and 20)

Numerator: Number responses 'Usually' or 'Always' that provider seemed to know important history, follow-up to give results from tests, and talked about all prescription medicines

Denominator: All responses



Children's Access to Primary Care - 12 to 24 months

Numerator: Number of children who had a visit with a primary care provider during the measurement period.

Denominator: Number of children ages 12 to 24 months as of June 30 MY

Children's Access to Primary Care - 25 months to 6 years

Numerator: Number of children who had a visit with a primary care provider during the measurement period.

Denominator: Number of children ages 25 months to 6 years as of June 30 MY

Children's Access to Primary Care - 7 to 11 years

Numerator: Number of children who had a visit with a primary care provider during the measurement period.

Denominator: Number of children ages 7 to 11 years as of June 30 MY

Children's Access to Primary Care - 12 to 19 years

Numerator: Number of children who had a visit with a primary care provider during the measurement period.

Denominator: Number of children ages 12 to 19 years as of June 30 MY

Controlling High Blood Pressure

Numerator: Number of people whose blood pressure was adequately controlled as follows:

- below 140/90 if ages 18-59;
- below 140/90 for ages 60 to 85 with a diabetes diagnosis; or
- below 150/90 ages 60 to 85 without a diagnosis of diabetes

Denominator: Number of people ages 18 to 85, who have hypertension

Diabetes Monitoring for People with Diabetes and Schizophrenia

Numerator: Number of people who had both an LDL-C test and an HbA1c test during the Measurement Year

Denominator: Number of people ages 18 to 64 with schizophrenia and diabetes

Diabetes Screening for people with Schizophrenia or Bipolar Disease who are using Antipsychotic Medication

Numerator: Number of people who had a diabetes screening test during the Measurement Year

Denominator: Number of people ages 18 to 64 years with Schizophrenia or Bipolar Disorder who were dispensed an antipsychotic medication

Discussion of Risks and Benefits of Aspirin Use

Numerator: Number of respondents who discussed risks and benefits of using aspirin with a doctor or health provider

Denominator: Number of respondents who are men, ages 46 to 79 years old and women who are 56 to 79 years old

ED Use by Uninsured

Numerator: The percentage of ED visits which are self-pay payer typology only for all hospitals in the PPS network in the Measurement Year

Denominator: The percentage of ED visits for self-pay payer typology only for all hospitals in the PPS network in the baseline Measurement Year

Engagement of Alcohol or Other Drug Dependence Treatment (Initiation and 2 visits within 44 days)

Numerator: Number of people who initiated treatment AND who has two or more additional services with a diagnosis of AOD within 30 days of the initiation visit

Denominator: Number of people age 13 and older with a new episode of AOD dependence



Flu Shots for Adults Ages 18 – 64

Numerator: Number of respondents who have had a flu shot

Denominator: Number of respondents, ages 18 to 64 years

Follow-up After Hospitalization for Mental Illness – within 7 days

Numerator: Number of discharges where the patient was seen on an ambulatory basis or who was in intermediate treatment with a mental health provider within 7 days of discharge

Denominator: Number of discharges between the start of the measurement period to 30 days before the end of the measurement period for patients ages 6 years and older who were hospitalized for treatment of selected mental health disorders

Follow-up After Hospitalization for Mental Illness – within 30 Days

Numerator: Number of discharges where the patient was seen on an ambulatory basis or who was in intermediate treatment with a mental health provider within 30 days of discharge

Denominator: Number of discharges between the start of the measurement period to 30 days before the end of the measurement period for patients ages 6 years and older who were hospitalized for treatment of selected mental health disorders

Follow-up Care for Children prescribed ADHD Medications - Initiation Phase

Numerator: Number of children who had one follow-up visit with a practitioner within the 30 days after starting the medication

Denominator: Number of children ages 6 to 12 years who were newly prescribed ADHD medication.

Follow-up Care for Children Prescribed ADHD Medications - Continuation Phase

Numerator: Number of children who in addition to the visit in the initiation phase had at least 2 follow-up visits in the 9-month period after the initiation phase ended

Denominator: Number of children ages 6 to 12 years old who were newly prescribed ADHD medication and remained on the medication for 7 months

Getting Timely Appointments, Care and Information (Q6, Q8 and Q10)

Numerator: Number of responses "Usually" or "Always" got appointment for urgent care or routine care as soon as needed, got answers the same day if called during the day

Denominator: Number who answered they called for appointments or called for information

HCAHPS Care Transition Metrics (Q23, Q24 and Q25)

Numerator: Average of hospital specific results for the Care Transition composite using "Strongly Agree" and "Agree" Responses

Denominator: Hospitals with HCAHPS participating in the PPS network

Health Literacy – Instructions Easy to Understand – (QHL13)

Numerator: Number responses 'Usually' or 'Always' that instructions for caring for condition were easy to understand

Denominator: Number who answered they saw provider for an illness or condition and were given instructions

Health Literacy – Describing How to Follow Instructions – (QHL14)

Numerator: Number responses 'Usually' or 'Always' that provider asked patient to describe how the instruction would be followed

Denominator: Number of respondents who answered they saw provider for an illness or condition and were given instructions



Health Literacy – Explained What to do if Illness Got Worse – (QHL16)

Numerator: Number responses 'Usually' or 'Always' that provider explained what to do if illness/condition got worse or came back

Denominator: Number who answered they saw provider for an illness or condition

Initiation of Alcohol & Other Drug Dependence Treatment (1 visit w/in 14 days)

Numerator: Number of people who initiated treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the index episode

Denominator: Number of people age 13 and older with a new episode of AOD dependence

Medical Assistance with Smoking and Tobacco Use Cessation – Advised to Quit

Numerator: Number of responses "Sometimes", "Usually" or "Always" were advised to quit

Denominator: Number of respondents, ages 18 and older, who smoke or use tobacco some days or every day

Medical Assistance with Smoking and Tobacco Use Cessation – Discussed Cessation Medication

Numerator: Number of responses "Sometimes", "Usually" or "Always" discussed cessation Medications

Denominator: Number of respondents, ages 18 and older who smoke or use tobacco some days or every day

Medical Assistance with Smoking and Tobacco Use Cessation – Discussed Cessation Strategies

Numerator: Number of responses "Sometimes", "Usually" or "Always" discussed cessation methods or strategies

Denominator: Number of respondents, ages 18 and older who smoke or use tobacco some days or every day

Non-use of Primary Care and Preventive Care Services

Numerator: The percentage of Medicaid members who do not have at least one claim with a preventive services CPT or equivalent code in the Measurement Year

Denominator: The percentage of Medicaid members who do not have at least one claim with a preventive services CPT or equivalent code in the baseline Measurement Year

Pediatric Quality Indicators (PDI) 90- Composite of All Measures

Numerator: Number of admissions which were in the numerator of one of the pediatric prevention quality indicators

Denominator: Number of people 6 to 17 years as of June 30 MY

Potentially Preventable Emergency Department Visits (for persons with BH diagnosis)

Numerator: Number of preventable emergency visits as defined by revenue and CPT codes

Denominator: Number of people with a BH diagnosis (excludes those born during the Measurement Year) as of June 30 of Measurement Year

Potentially Preventable Emergency Room Visits

Numerator: Number of preventable emergency visits as defined by revenue and CPT codes

Denominator: Number of people (excludes those born during the Measurement Year) as of June 30 of Measurement Year

Potentially Avoidable Readmissions

Numerator: Number of readmissions chains (at risk admissions followed by one or more clinically related readmissions within 30 days of discharge)

Denominator: Number of people as of June 30 of the Measurement Year



Prevention Quality Indicator # 7 (Hypertension)

Numerator: Number of admissions with a principal diagnosis of hypertension

Denominator: Number of people 18 years and older who were enrolled in Medicaid for at least one month as of June 30 of the Measurement Year

Prevention Quality Indicator # 8 (Heart Failure)

Numerator: Number of admissions with a principal diagnosis of heart failure

Denominator: Number of people 18 years and older who were enrolled in Medicaid for at least one month as of June 30 of the Measurement Year

Primary Care - Length of Relationship – Q3

Numerator: Percentage of responses for at least 1 year or longer

Denominator: All Responses

Primary Care - Usual Source of Care – Q2

Numerator: Percent of Responses "Yes"

Denominator: All Responses

Prevention Quality Indicator (PQI) - 90 Composite of all Measures

Numerator: Number of admissions which were in the numerator of one of the adult prevention quality indicators

Denominator: Number of people 18 years and older who were enrolled in Medicaid for at least one month as of June 30 of the Measurement Year

Screening for Clinical Depression and Follow-Up

Numerator: Number of people screened for clinical depression using a standardized depression screening tool and if positive, with follow-up within 30 days

Denominator: Number of people with a qualifying outpatient visit who are 18 and older

Statin Therapy for Patients with Cardiovascular Disease - Received Statin Therapy

Numerator: Number of people who were dispensed at least one high or moderate-intensity statin medication

Denominator: Number of males age 21 to 75 or females age 40 to 75 who have had an MI, CABG or PCI in the year prior or a diagnosis of ischemic vascular disease in both the Measurement Year and year prior

Statin Therapy for Patients with Cardiovascular Disease - Statin Adherence 80%

Numerator: Number of people who achieved a proportion of days covered of 80% for the treatment period

Denominator: Number of males age 21 to 75 or females age 40 to 75 who have had an MI, CABG or PCI in the year prior or a diagnosis of ischemic vascular disease in both the Measurement Year and year prior AND who were dispensed at least one high or moderate-intensity statin medication